



⑪ Publication number : **0 594 285 A2**

⑫ **EUROPEAN PATENT APPLICATION**

⑲ Application number : **93306526.0**

⑥ Int. Cl.⁵ : **C08G 77/388, C07F 7/10**

⑳ Date of filing : **18.08.93**

③ Priority : **21.08.92 US 934051**

④ Date of publication of application :
27.04.94 Bulletin 94/17

⑧ Designated Contracting States :
DE FR GB

⑦ Applicant : **GENERAL ELECTRIC COMPANY**
1 River Road
Schenectady, NY 12345 (US)

⑦ Inventor : **Lucarelli, Michael Anthony**
7 McGinnis Place
Mattoon, Illinois 61938 (US)
Inventor : **Raleigh, William John**
10 Glen Street
Rensselaer, New York 12144 (US)

⑦ Representative : **Pratt, Richard Wilson et al**
London Patent Operation G.E. Technical
Services Co. Inc. Essex House 12/13 Essex
Street
London WC2R 3AA (GB)

⑤ Amino acid functionalized silicones.

⑤ Organofunctional siloxanes useful in a wide variety of personal care and plastics applications are disclosed having a carboxyalkylaminoalkyl functionality.

EP 0 594 285 A2

The present invention relates to silicones useful in both the personal care and plastics industries. More particularly, the present invention relates to an organofunctional silicone. Most particularly, the present invention relates to amino acid functionalized silicones.

5 BACKGROUND OF THE PRESENT INVENTION

Organofunctional silicones are well known in the art. The siloxane may be functionalized with substituents such as carboxy, amido, ketoxy, epoxy, vinyl, hydride, mercapto, amino, glycidoxo, methacryloxy and alkoxy groups. These organofunctionalities are generally employed to obtain specific physical and chemical properties. Aminoalkyl-substituted silicones, for example, increase water solubility and are used in automobile polishes to enhance durability and confer detergent resistance through bonding with paint films. Other functionalized fluids find use in applications such as textile lubricants, softeners and antistats. A variety of amino functional siloxanes are described in the prior art.

Brumbill, United States Patent No. 4,785,067, teaches a protective coating consisting essentially of a polysiloxane copolymer prepared by the reaction of a trimethyl end blocked polyamine silicone fluid including amino alkyl groups with a saturated aliphatic carboxylic acid groups to produce a polysiloxane copolymer including alkyl amine groups.

van Hooijdonk, United States Patent No. 5,028,485, teaches a pressure sensitive adhesive composition suitable for skin contact comprising a silane compound having at least one amino functional group.

Imperante et al., United States Patent No. 5,115,049, teach the preparation of fatty carboxylic acid salts of organofunctional silicone amines, where the amino pendant functionality is present within the polymer and which are produced by the neutralization of a silicone amine with a fatty carboxylic acid.

It has now been found that novel amino acid functionalized silicones can be produced, and that such compounds are useful in a variety of plastic and personal care applications.

25 SUMMARY OF THE INVENTION

According to the present invention there is provided an organofunctional silicone composition comprising those of the formula: (i) $TD_xM'_3$; (ii) $TD_xD'_yM'_3$; (iii) $TD'_yM'_3$; (iv) $TD_xD'_yM_3$; (v) TD'_yM_3 ; (vi) $M'D_xD'_yM'$; (vii) $M'D_xM'$; (viii) MD'_yM ; (ix) $MD_xD'_yM$; (x) $M'D'_yM'$; (xi) $M'Q$; (xii) $(D')_z$; or (xiii) a combination of any of the foregoing; wherein T represents a trifunctional siloxy group of the formula $RSiO_{3/2}$ wherein R represents a saturated or unsaturated monovalent hydrocarbon radical; D represents a difunctional siloxy group of the formula $R_2SiO_{2/2}$ wherein each R is independently defined as above; D' represents a difunctional siloxy group of the formula $RR'SiO_{2/2}$ wherein each R is independently defined as above, R' is a carboxyalkyl aminoalkyl group of the formula $(HO_2C)(CH_2)_mN(R^2)(CH_2)_n$ where R^2 represents hydrogen or an alkyl group of from 1 to 10 carbon atoms, and m and n each independently vary from 1 to 10; M represents a monofunctional siloxy group of the formula $R_3SiO_{1/2}$ wherein each R is independently defined as above; M' represents a monofunctional siloxy group of the formula $R_2R'SiO_{1/2}$ wherein each R is independently defined as above and R' is as above defined; Q represents a quadrifunctional siloxy group of the formula $SiO_{4/2}$; x is greater than 1, y is greater than 1 and z is equal to or greater than 3.


Also according to the present invention there is provided a process for preparing the amino acid functionalized silicones of the present invention comprising the steps of: (a) hydrosilating a silicone hydride compound with a lactam to form an amide functionalized silicone; and (b) hydrolyzing the amide functionalized silicone of step (a) to produce an amino acid functionalized silicone.

45 DETAILED DESCRIPTION OF THE PRESENT INVENTION

The novel amino acid functionalized silicones of the present invention are generally prepared by hydrosilating the appropriate linear, branched or cyclic silicone hydride with a lactam and then hydrolyzing the lactam functionalized silicone to produce the amino acid functionalized silicones of the present invention.

The silicone hydrides for use in the practice of the present invention are typically those which correspond to the product amino acid functionalized siloxanes, and are generally selected from $TD_xM'_3$; $TD_xD'_yM'_3$; $TD'_yM'_3$; $TD_xD'_yM_3$; TD'_yM_3 ; $M'D_xD'_yM'$; $M'D_xM'$; MD'_yM ; $MD_xD'_yM$; $M'D'_yM'$; $M'Q$; $(D')_z$; or combinations of any of the foregoing where M' is a monofunctional siloxy hydride of the formula $(R)_2(H)SiO_{1/2}$; D is a difunctional siloxy hydride of the formula $(R)(H)SiO_{2/2}$ and T, D, M, Q, R, x, y, and z are as defined hereinabove. The siloxy hydrides are available commercially and can be prepared by methods known to those skilled in the art.

The lactams useful in the present invention for the hydrosilation reaction are generally of the formula



wherein R^3 is an alkenyl group of from 2 to about 10 carbon atoms and q is an integer ranging from 1 to about 10. A preferred lactam for use in the practice of the present invention is *n*-vinyl pyrrolidone, wherein R^3 is vinyl, and q is 2 in the above formula.

The lactam is generally reacted with the appropriate silicone hydride by hydrosilation techniques well known to those of ordinary skill in the art. In general the hydrosilation reaction proceeds in the presence of a Group VIII precious metal complex, i.e. platinum, rhodium and/or ruthenium complexes. Such catalyst complexes are well described in the patent literature, e.g., Karstedt, United States Patent No. 3,775,452; Ashby et al., United States patent No. 4,288,345; Bailey et al., United States Patent No. 3,336,239; Ashby, United States Patent No. 4,421,903; Lamoreaux, United States Patent No. 3,220,972; and Lewis, United States Patent No. 4,946,818. See also, J. Organometallic Chem. 408 (1991) pp. 297-304. A preferred catalyst for use in the present invention comprises tris(dibutylsulfide) rhodium trichloride.




The lactam modified siloxanes are then reacted in a hydrolysis reaction according to techniques known to those skilled in the art. Typically, the hydrolysis reaction is conducted in the presence of an acid, such as dilute sulfuric acid. However, other acids may be employed. The hydrolysis reaction acts to open the amine ring structure to give the corresponding acid. The product is an amino acid functionalized siloxane.

The organofunctional siloxanes of the present invention may be branched (T or Q), linear (D) or cyclic ($[(D)_z]$). They are of the general formulae: (i) $\text{TD}_x\text{M}'_3$; (ii) $\text{TD}_x\text{D}'_y\text{M}'_3$; (iii) $\text{TD}'_y\text{M}'_3$; (iv) $\text{TD}_x\text{D}'_y\text{M}_3$; (v) $\text{TD}'_y\text{M}_3$; (vi) $\text{M}'\text{D}_x\text{D}'_y\text{M}'$; (vii) $\text{M}'\text{D}_x\text{M}'$; (viii) $\text{MD}'_y\text{M}$; (ix) $\text{MD}_x\text{D}'_y\text{M}$; (x) $\text{M}'\text{D}'_y\text{M}'$; (xi) $\text{M}'\text{Q}$; (xii) $(\text{D}')_z$; or (xiii) a combination of any of the foregoing. In preferred embodiments, x and y are each independently greater than about 1, preferably each independently ranges from about 1 to about 1000, and more preferably each ranges from about 1 to about 100. In the case of cyclics, z is typically equal to or greater than about 3 and preferably varies from about 4 to about 12.

In the above formulae, each R independently represents a saturated or unsaturated monovalent hydrocarbon, typically containing no more than six carbon atoms and selected from those such as alkyl radicals, e.g., methyl, ethyl and isopropyl; cycloaliphatic radicals, e.g., cyclopentyl and cyclohexenyl; olefinic radicals, e.g., vinyl and allyl; and the phenyl radical.

R^1 represents a carboxyalkyl aminoalkyl group of the formula $(\text{HO}_2\text{C})(\text{CH}_2)_m\text{N}(\text{R}^2)(\text{CH}_2)_n$, where R^2 represents hydrogen or an alkyl group of from 1 to 10 carbon atoms, and m and n each independently vary from 1 to 10. In preferred embodiments, R^1 is selected from N-(3-carboxypropyl)-aminoethyl; N-(3-carboxybutyl)-aminoethyl; N-(3-carboxymethyl)aminopentyl and the like. Particularly useful in the practice of the present invention is N-(3-carboxypropyl)aminoethyl.

The following reaction scheme depicts the general reaction route for linear materials reacted with *n*-vinyl pyrrolidone.



25

30

35

EXAMPLE 1

45

50

55

55

a vacuum pump to isolate the n-vinyl pyrrolidone adduct. 32.5 g of oil was isolated. IR 1684 cm^{-1} ; NMR CDCl_3 standard: 0.05 ppm s,s 12H (Me-Si), 1.09 m 4H ($\text{CH}_2\text{-Si}$), 2.00 m 4H (CH_2), 2.25 m 4H (CH_2), 3.27 m 4H ($\text{CH}_2\text{-N}$), 3.41 m 4H ($\text{CH}_2\text{-N}$). The NMR assignments are based on the alpha-alpha adduct and comparison of spectra of various substituted pyrrolidones. See, Bovey, F.A., "NMR Data Tables for Organic Compounds", Interscience Publishers, 1967, V.1.

To the remainder of the mixture, was added 40 ml of water which contained 4 drops of sulfuric acid. The mixture was heated at 90°C for one hour and then allowed to cool. The mixture was then extracted with toluene, the organic layer subsequently being dried over sodium sulfate and rotoevaporated to afford 67 g of an amino acid functionalized siloxane oil of the present invention. IR 1678 cm^{-1} broad. NMR CDCl_3 standard: 0.00 ppm s,s 12H ($\text{CH}_3\text{-Si}$); 1.02 m 4H ($\text{CH}_2\text{-Si}$), 1.82 m 4H, 2.19 m 4H, 3.25 m 4H and 3.36 m 4H.

The above-mentioned patents and publications are all hereby incorporated by reference.

Many variations of the present invention will suggest themselves to those skilled in the art in light of the above-detailed description. Other siloxanes may be employed in the present invention, including $\text{TD}_x\text{D}'_y\text{M}'_z$, $\text{TD}'_y\text{M}'_3$, $\text{TD}_x\text{D}'_y\text{M}_3$, $\text{TD}'_y\text{M}_3$, $\text{M}'\text{D}_x\text{M}'$, $\text{MD}'_y\text{M}$, $\text{MD}_x\text{D}'_y\text{M}$, $\text{M}'\text{D}'_y\text{M}'$, $\text{M}'\text{Q}$ and $(\text{D}')_z$ may all be employed in the present invention. Further, a wide variety of lactams other than n-vinyl pyrrolidone are contemplated for use in the present invention. All such obvious modifications are within the full intended scope of the appended claims.

Claims

1. An organofunctional silicone composition comprising those of the formula:

- (i) $\text{TD}_x\text{M}'_3$;
- (ii) $\text{TD}_x\text{D}'_y\text{M}'_3$;
- (iii) $\text{TD}'_y\text{M}'_3$;
- (iv) $\text{TD}_x\text{D}'_y\text{M}_3$;
- (v) $\text{TD}'_y\text{M}_3$;
- (vi) $\text{M}'\text{D}_x\text{D}'_y\text{M}'$;
- (vii) $\text{M}'\text{D}_x\text{M}'$;
- (viii) $\text{MD}'_y\text{M}'$;
- (ix) $\text{MD}_x\text{D}'_y\text{M}$;
- (x) $\text{M}'\text{D}'_y\text{M}'$;
- (xi) $\text{M}'\text{Q}$;
- (xii) $(\text{D}')_z$; or

(xiii) a combination of any of the foregoing wherein T represents a trifunctional siloxy group of the formula $\text{RSiO}_{3/2}$ wherein R represents independently a saturated or unsaturated monovalent hydrocarbon radical; D represents a difunctional siloxy group of the formula $\text{R}_2\text{SiO}_{2/2}$ wherein each R is independently defined as above; D' represents a difunctional siloxy group of the formula $\text{RR}'\text{SiO}_{2/2}$ wherein each R is independently defined as above, R' represents a carboxyalkyl aminoalkyl group of the formula $(\text{HO}_2\text{C})(\text{CH}_2)_m\text{N}(\text{R}^2)(\text{CH}_2)_n$ where R^2 represents hydrogen or an alkyl group of from 1 to about 10 carbon atoms, and m and n each independently vary from 1 to about 10; M represents a monofunctional siloxy group of the formula $\text{R}_3\text{SiO}_{1/2}$ wherein each R is independently defined as above; M' represents a monofunctional siloxy group of the formula $\text{R}_2\text{R}'\text{SiO}_{1/2}$ wherein each R is independently defined as above and R' is as above defined; Q represents a quadrifunctional siloxy of the formula $\text{SiO}_{4/2}$; x is greater than 1, y is greater than 1 and z is equal to or greater than 3.

2. An organofunctional silicone composition as defined in Claim 1 wherein R is methyl.

3. An organofunctional silicone composition as defined in Claim 1 or Claim 2 wherein each R^2 is independently ethyl or hydrogen.

4. An organofunctional silicone composition as defined in any preceding claim wherein x ranges from 1 to about 1000.

5. An organofunctional silicone composition as defined in any preceding claim wherein y ranges from 1 to about 1000.

6. A method of preparing an organofunctional silicone composition comprising units of the formula
(i) $\text{TD}_x\text{M}'_3$;

- (ii) $TD_xD'_yM'_z$;
 (iii) $TD'_yM'_z$;
 (iv) $TD_xD'_yM_3$;
 (v) TD'_yM_3 ;
 5 (vi) $M'D_xD'_yM'$;
 (vii) $M'D_xM'$;
 (viii) MD'_yM ;
 (ix) $MD_xD'_yM$;
 (x) $M'D'_yM'$;
 10 (xi) $M'Q$;
 (xii) $(D')_z$; or
 (xiii) a combination of any of the foregoing wherein T represents a trifunctional siloxy group of the formula $RSiO_{3/2}$ wherein R represents independently a saturated or unsaturated monovalent hydrocarbon radical; D represents a difunctional siloxy group of the formula $R_2SiO_{2/2}$ wherein each R is independently defined as above; D' represents a difunctional siloxy group of the formula $RR'SiO_{2/2}$ wherein each R is independently defined as above, R' represents a carboxyalkyl aminoalkyl group of the formula $(HO_2C)(CH_2)_mN(R^2)(CH_2)_n$ where R^2 represents hydrogen or an alkyl group of from 1 to about 10 carbon atoms, and m and n each independently vary from 1 to about 10; M represents a monofunctional siloxy group of the formula $R_3SiO_{1/2}$ wherein each R is independently defined as above; M' represents a monofunctional siloxy group of the formula $R_2R'SiO_{1/2}$ wherein each R is independently defined as above and R' is as above defined; Q represents a quadrifunctional siloxy of the formula $SiO_{4/2}$; x is greater than 1, y is greater than 1 and z is equal to or greater than 3;
 comprising the steps of:
 (a) hydrosilating a silicone hydride compound with a lactam to form an amide functionalized silicone;
 25 and
 (b) hydrolyzing the amide functionalized silicone of step (a) to produce an amino acid functionalized silicone.

7. A process as defined in Claim 6 wherein said silicone hydride is selected from $TD_xM''_3$; $TD_xD'_yM''_3$;
 30 $TD'_yM''_3$; $TD_xD'_yM_3$; TD'_yM_3 ; $M''D_xD'_yM''$; $M''D_xM''$; MD'_yM ; $MD_xD'_yM$; $M''D'_yM''$; $M''Q$; $(D'')_z$; or combinations of any of the foregoing where M'' is a monofunctional siloxy hydride of the formula $(R)_2(H)SiO_{1/2}$; D is a difunctional siloxy hydride of the formula $(R)(H)SiO_{2/2}$ and T, D, M, Q, R, x, y, and z are as defined above.
8. A process as defined in Claim 6 or Claim 7 wherein said lactam has the general formula



wherein R^3 is an alkenyl group of from 2 to 10 carbon atoms and q is an integer ranging from 1 to about 10.

9. A process as defined in any one of Claims 6 to 8 wherein the hydrosilating step (a) is carried out with a catalyst selected from a rhodium catalyst or a platinum catalyst.
10. A process as defined in any one of Claims 6 to 9 wherein the hydrolyzing step (b) is carried out with a dilute sulfuric acid.